

Atty. Docket No.  
A35478 066123.0125

Serial No.  
10/687,402

**INFORMATION DISCLOSURE STATEMENT  
BY APPLICANT**

(Use several sheets if necessary)

Applicant  
Tyagi et al.

Filing Date  
October 16, 2003

Group  
1651

**U.S. PATENT DOCUMENTS**

*Exam. Init.	Document No.	Date	Name	Class	Subclass	Filing Date if Appropriate
JS	6 1 6 2 6 2 7	12/19/2000	Inouye et al.	435	194	

**FOREIGN PATENT DOCUMENT**

Document No.	Date	Country	Class	SubClass	Translation Yes No
0 3 0 6 6 8 3 8	08/14/2003	WIPO			

**OTHER DOCUMENTS (including Author, Title Date, Pertinent Pages, Etc.)**

JS	Anderson D.H., Harth G., Horwitz M.A. and Eisenberg D. (2001) An interfacial mechanism and a class of inhibitors inferred from two crystal structures of the <i>Mycobacterium tuberculosis</i> 30 kDa major secretory protein (Antigen 85B) a mycolyl transferase. <i>J. Mol. Biol.</i> 307, 671-681
	Armitage, L.Y., Jagannath, C., Wanger, A.R. and Norris, S.J. (2000) Disruption of gene encoding antigen 85A and 85B of <i>Mycobacterium tuberculosis</i> H37Rv: Effect on growth in culture and in macrophages. <i>Infect. Immun.</i> 68, 76-78
	Baikalov, I., Schroder, I., Kaczor - Grzeskowiak, M., Grzeskowiak, K., Gunsalus, R.P. and Dickerson, R.E. (1996) Structure of the <i>Escherichia coli</i> response regulator NarL. <i>Biochemistry</i> 35, 11053-11061
	Baltch, A.L., Smith, R.P., Ritz, W.J. and Bopp, L.H. (1998) Comparison of inhibitory and bactericidal activities and postantibiotic effects of LY333328 and Ampicillin used singly and in combination against vancomycin - resistant <i>Enterococcus faecium</i> . <i>Antimicrob. Agents Chemother.</i> 42, 2564-2568
	Barrett, J.F. and Hoch, J.A. (1998) Two-component signal transduction as a target for microbial anti - infective therapy. <i>Antimicrob. Agents Chemother.</i> 42, 1529-1536
	Barry, C.E. III, Slayden, R.A., Simpson, A.E. and Lee, R.E. (2000) Use of genomics and combinatorial chemistry in the development of new antimycobacterial drugs. <i>Biochem. Pharmacol.</i> 59, 221-231
↓	Boon, C., Li, R., Qi, R. and Dick, T. (2001) Proteins of <i>Mycobacterium bovis</i> BCG induced in Wayne dormancy model. <i>J. Bacteriol.</i> 182, 2672-2676

NY02:505269.1

1

Examiner

Date Considered

12-20-04

\* Examiner: Initial citation considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

**INFORMATION DISCLOSURE STATEMENT  
BY APPLICANT**  
(Use several sheets if necessary)

Applicant  
Tyagi et al.

Filing Date  
October 16, 2003


Group  
1651

✓		Bradford, M. M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein - dye binding. <i>Anal. Biochem.</i> 72, 248 - 254
		Chen, P., Ruiz, R.E., Li, Q., Silver, R.F. and Bishai, W.R. (2000) Construction and characterization of <i>M. tuberculosis</i> mutant lacking the alternate sigma factor, <i>sigF</i> . <i>Infect. Immun.</i> 68, 5575 - 5580
		Cole et al. (1998) Deciphering the biology of <i>Mycobacterium tuberculosis</i> from the complete genome sequence. <i>Nature</i> 393, 537 - 544
		Collins D.M., Kawakami R.P., de Lisle G.W., Pascopella L., Bloom B.R. and Jacobs W.R. Jr. (1995) Mutation of the principal sigma factor causes loss of virulence in a strain of the <i>Mycobacterium tuberculosis</i> complex. <i>Proc. Natl. Acad. Sci. USA</i> 92, 8036 - 8040
		Cooper J.B., McIntyre K., Badasso M.O., Wood S.P., Zhang Y., Garbe T.R. and Young, D. (1995) X-ray structure analysis of the iron-dependent superoxide dismutase from <i>Mycobacterium tuberculosis</i> at 2.0 Angstroms resolution reveals novel dimer-dimer interactions. <i>J. Mol. Biol.</i> 246, 531 - 544
		Cox J.S., Chen B., McNeil M. and Jacobs W.R. Jr. (1999) Complex lipid determines tissue-specific replication of <i>Mycobacterium tuberculosis</i> in mice, <i>Nature</i> 402, 79 - 83
		Dasgupta, N., Kapur, V., Singh, K.K., Das, T.K., Sachdeva, S., Jyothisri, K. and Tyagi, J.S. (2000) Characterization of a two-component system, <i>devR-devS</i> , of <i>Mycobacterium tuberculosis</i> . <i>Tuber. Lung Dis.</i> 80, 141 - 159
		DeMaio J., Zhang Y., Ko C., Young D.B. and Bishai W.R. (1996) A stationary-phase stress-response sigma factor from <i>Mycobacterium tuberculosis</i> . <i>Proc. Natl. Acad. Sci. (USA)</i> 93, 2790-2794
		Denis et al. (1999) Synthesis and antibacterial activity of HMR 3647, a new ketolide highly potent against erythromycin - resistant and susceptible pathogens. <i>Bioorg. Med. Chem. Lett.</i> 9, 3075 - 3080
		Deschenes, R.J., Lin, H., Ault, A.D. and Fassler, J.S. (1999) Antifungal properties and target evaluation of three putative bacterial histidine kinase inhibitors. <i>Antimicrob. Agents Chemother.</i> 43, 1700 - 1703
		Domagala, J.M., Alessi, D., Cummings, M., et al. (1998) Bacterial two-component signalling as a therapeutic target in drug design: Inhibition of NRII by diphenolic methanes (bisphenols). <i>Adv. Exp. Med. Biol.</i> 456, 269 - 286
✓		Doukhan, L., Predich, M., Nair, G., Dussurget, O., Mandic-Mulec, I., Cole, S.T., Smith, D.R. and Smith, I. (1995) Genomic organization of the mycobacterial sigma gene cluster. <i>Gene</i> 165, 67 - 70

NY02:505269.1

2

Examiner



Date Considered

12 - 20 - 04

**INFORMATION DISCLOSURE STATEMENT  
BY APPLICANT**

(Use several sheets if necessary)

Applicant  
Tyagi et al.

Filing Date  
October 16, 2003

Group  
1651

28		Dziejman, M. and Mekalanos, J.J. (1995) two-component signal transduction and its role in the expression of bacterial virulence factors in Hock, J.A. and Silhavy, T.J. Eds. Two component signal transduction. Washington, DC. American Society for Microbiology. Pp. 305-317
		El-Masry, A.H., Fahony, H.H. and Abdelwahed, S.H.A. (2000) Synthesis and anti microbial activity of some new benzimidazole derivatives. <i>Molecules</i> 5, 1429 - 1438
		Frechette, R.F., Beach, M.J., Bernstein, J. et al. (1997) Novel benzoxazine derivatives with inhibitory activity against bacterial two-component signal transduction systems. Abstract No. 152, Book of Abstracts, 214th ACS National Meeting, Las Vegas, NV, USA
		Glickman, M.S., Cox, J.S. and Jacobs, W.R. Jr. (2000) A novel mycolic acid cyclopropane synthetase is required for coding, persistence and virulence of <i>Mycobacterium tuberculosis</i> . <i>Mol. Cell</i> 5, 717 - 727
		Grange, J.M. (1992) The mystery of the mycobacterial 'persistor'. <i>Tuber. Lung Dis.</i> 73, 249 - 251..
		Haydel, S.E., Dunlap, N.E. and Benjamin, W.H. Jr. (1999) <i>In vitro</i> evidence of two-component system phosphorylation between the <i>Mycobacterium tuberculosis</i> TrcR/ TrcS proteins <i>Micro. Patho.</i> 26 195-206
		Himpens, S., Loch, C. and Supply, P. (2000) Molecular characterization of the mycobacterial SenX3 - RegX3 two-component system: evidence for autoregulation. <i>Microbiol.</i> 146, 3091 - 3098
		Jackson, M., Crick, D.C. and Brennan, P.J. (2000) Phosphatidylinositol is an essential phospholipid of mycobacteria <i>J. Biol. Chem.</i> 275, 30092 - 30099
		Kinger, A.K. and Tyagi, J.S. (1993) Identification and cloning of genes differentially expressed in the virulent strain of <i>Mycobacterium tuberculosis</i> . <i>Gene</i> 131, 113 - 117
		Klimesova, V., Koc, J., Pour, M., Stachel, J., Waisser, K. and Kaustova, J. (2002a) Synthesis and preliminary evaluation of benzimidazole derivatives as antimicrobial agents. <i>Eur. J. Med. Chem.</i> 37, 409 - 418
		Klimesova, V., Koc, J., Waisser, K. and Kaustova, J. (2002b) New benzimidazole derivatives as antimycobacterial agents. <i>Il Farmaco</i> , 57, 259 - 265
		Kramer, MJ and Grunberg, E. (1973) Effect of ethidium bromide against transplantable tumors in mice and rats. <i>Chemotherapy</i> , 19, 254 - 258

NY02:505269.1

3

Examiner



Date Considered

12-20-04

**INFORMATION DISCLOSURE STATEMENT  
BY APPLICANT**  
(Use several sheets if necessary)

Applicant  
Tyagi et al.

Filing Date  
October 16, 2003

Group  
1651

87		Phetsuksiri, B., Baulard, A.R., Cooper, A.M., Minnikin, D.E., Douglas, J.D., Besra, G.S. and Brennan, P.J. (1999) Antimycobacterial activities of isoxyl and new derivatives through the inhibition of mycolic acid synthesis. <i>Antimicrob. Chemother.</i> 43, 1042-1051
		Li, Z., Kelley, C., Collins, F., Rouse, D. and Morris, S. (1998) Expression of katG in <i>Mycobacterium tuberculosis</i> is associated with its growth and persistence in mice and guinea pigs. <i>J. Infect. Dis.</i> 177, 1030 - 1035
		Macielag, M.J. and Goldschmidt, R. (2000) Inhibitors of bacterial two-component signalling system. <i>Exp. Opin. Invest. Drugs</i> 9, 2351 - 2369
		Manganelli, R., Voskuil, M.I., Schoolnik, G.K. and Smith, I. (2001) The <i>Mycobacterium tuberculosis</i> ECF sigma factor sigmaE: role in global gene expression and survival in macrophages. <i>Mol. Microbiol.</i> 41, 423-437
		Matsushita, M. and Janda, K.D. (2002) Histidine kinases as targets for new antimicrobial agents. <i>Bioor. Med. Chem.</i> 10, 855 - 867
		Mayuri, Bagchi, G., Das, T.K. and Tyagi, J.S. (2002) Molecular analysis of the dormancy response in <i>Mycobacterium smegmatis</i> . Expression analysis of the genes encoding DevR - DevS two-component system, Rv3134c and chaperone - crystalline homologues. <i>FEMS Microbiol. Lett.</i> 211, 231 - 237
		McKinney, J.D., Bentrup, K.H., Munoz-Elias, E.J., Miczak, A., Chen, B., Chen, W.T., Swenson, D., Sacchettini, J.C. Jacobs, W.R. Jr. and Russell, D.G.. (2000) Persistence of <i>M. tuberculosis</i> in macrophages and mice requires the glyoxylate shunt enzyme isocitrate lyase. <i>Nature</i> 406, 735 - 738
		Mitchison, D.A. (1998) How drug resistance emerges as a result of poor compliance during short - chemotherapy for tuberculosis. <i>Int. J. Tuberc. Lung Dis.</i> 2, 10 - 15
		Mukamolova G.V., Kaprelyants A.S., Young D.I., Young M. and Kell D.B. (1998) A bacterial cytokine. <i>Proc. Nat Acad. Sci. USA.</i> 95, 8916 - 8921
		Parkinson, J.S. and Kofoed, E.C. (1992) Communication modules in bacterial signalling proteins. <i>Ann. Rev. Genet.</i> 26, 71 - 112
		Parrish, N., Dick, J.D. and Bishai, W.R. (1998) Mechanisms of latency in <i>Mycobacterium tuberculosis</i> . <i>Trends Microbiol.</i> 6: 107 -112
		Perez, E., Samper, S., Bordas, Y., Guilhot, C., Gicquel, B. and Martin, C. (2001) An essential role for <i>phoP</i> in <i>Mycobacterium tuberculosis</i> virulence. <i>Mol. Micro.</i> 41, 179 - 87

NY02:505269.1

4

Examiner



Date Considered

12-20-04

**INFORMATION DISCLOSURE STATEMENT  
BY APPLICANT**

(Use several sheets if necessary)

Applicant  
Tyagi et al.

Filing Date  
October 16, 2003

Group  
1651

		Pohl, E., Holmes, R.K. and Hol, W.G. (1999) Crystal structure of the iron-dependent regulator (IdeR) from <i>Mycobacterium tuberculosis</i> shows both metal binding sites fully occupied. <i>J. Mol. Biol.</i> 285, 1145 - 1156
		Raman, S., Song, T., Puyang, X., Bardarov, S., Jacobs, W.R. Jr. and Husson, R.N. (2001) The alternative sigma factor SigH regulates major components of oxidative and heat stress responses in <i>Mycobacterium tuberculosis</i> . <i>J. Bacteriol.</i> 183, 6119 - 6125
		Ronning, D.R., Klabunde, T., Besra, G.S., Vissa, V.D., Belisle, J.T. and Sacchettini, J.C. (2000) Crystal structure of the secreted form of antigen 85C reveals potential targets for mycobacterial drugs and vaccines. <i>Nat. Struct. Biol.</i> 7 141 - 146
		Roychoudhary, S., Zielinski, N.A., Ninfa, A.J., Allen, N.E., Jungheim, L.N., Nicas, T.I. and Chakrabarty, A.M. (1993) Inhibitors of two-component signal transduction systems: Inhibition of alginate gene activation in <i>Pseudomonas aeruginosa</i> . <i>Proc. Natl. Acad. Sci. USA.</i> 90, 965 - 969
		Sherman, D.R., Voskuil, M., Schnappinger, D., Liao, R., Harrell, M.I. and Schoolnik, G.K. (2001) Regulation of the <i>Mycobacterium tuberculosis</i> hypoxia response gene encoding $\alpha$ -crystallin. <i>Proc. Natl. Acad. Sci. USA</i> 98, 7534 - 7539
		Stead, W.W. (1967) Pathogenesis of a first episode of chronic pulmonary tuberculosis in man: recrudescence of residuals of the primary infection or exogenous reinfection? <i>Am. Rev. Respir. Dis.</i> 95, 729 - 745
		Stead, W.W., Kerby, D.P., Schleuter, D.P. and Jordahl, C.W. (1968) The clinical spectrum of primary tuberculosis in adults. Confusion with reinfection in the pathogenesis of chronic tuberculosis. <i>Ann. Intern. Med.</i> 68, 731 - 745
		Stephenson, K., Yamaguchi, Y., Hoch, J.A. (2000) The mechanism of action of inhibitors of bacterial two-component signal transduction systems. <i>J. Biol. Chem.</i> 275, 38900 - 38904.
		Stock, A.M., Robinson, V.L. and Goudreau, P.N. (2000) Two-component signal transduction. <i>Ann. Rev. Biochem</i> 69, 183 - 215
		Stock, J.B., Surette, M.G., Levit, M. and Park, P. (1995) Two-component signal transduction systems: structure-function relationships and mechanisms of catalysis. In: Hock, J.A. and Silhavy, T.J. Eds. Two component signal transduction. Washington, DC. American Society for Microbiology. Pp. 25 - 51
		Stover, C.K., Warren, P., VanDevanter, D.R., Sherman, D.R., Arain, T.M., Langhorne, M.H., Anderson, S.W., Towell, J.A., Yuan, Y., McMurray, D.N., Kreiswirth, B.E., Barry, C.E. III and Baker, W.R. (2000) A small molecule nitroimidazole drug candidate for treatment of tuberculosis. <i>Nature</i> 405, 962 - 966
		Strauch, M.A., De Mendoza, D. and Hoch, J.A. (1992) cis - Unsaturated fatty acids specifically inhibit a signal - transducing protein kinase required for initiation of sporulation in <i>B. subtilis</i> . <i>Mol. Microbiol.</i> 6, 2909 - 2917

NY02:505269.1

5

Examiner

Date Considered



12-20-04

**INFORMATION DISCLOSURE STATEMENT  
BY APPLICANT**  
(Use several sheets if necessary)

Applicant  
Tyagi et al.

Filing Date  
October 16, 2003

Group  
1651

		Ulijasz, A.T. and Weisblum, B. (1999) Dissecting the VanRS signal transduction pathway with specific inhibitors. <i>J Bacteriol.</i> 181, 627 - 631
		Upton, A., Johnson, N., Sandy, J. and Sim, E. (2001) Arylamine N-acetyltransferases - of mice, men and microorganisms. <i>Trends Pharmacol. Sci.</i> 22, 140 - 146
		Urbanski, M.J., Xiang, M.A., Foleno, B.D. et al. (1997) Novel cyclohexene derivatives with histidine protein kinase inhibitory activity - potential new antibacterial agents. Abstract No. 270, Book of Abstracts, 214th ACS National Meeting, Las Vegas, NV, USA
		Wallis, N.G. (1999) Bacterial two-component signal transduction systems as drug targets. <i>Curr. Opin. Anti-infect. Invest. Drugs.</i> 1, 428 - 434
		Weber, I., Fritz, C., Rutkowski, S., Kreft, A. and Bange, F.-C. (2000) Anaerobic nitrate reductase ( <i>narGHJI</i> ) activity of <i>Mycobacterium bovis</i> BCG <i>in vitro</i> and its contribution to virulence in immunodeficient mice. <i>Mol. Microbiol.</i> 35 1017 - 1025
		West, M.L. and Fairlie, D.P. (1995) Targeting HIV-1 protease: a test of drug-design methodologies. <i>Trends Pharmacol. Sci.</i> 16, 67 - 94
		WHO report on the Global Tuberculosis Epidemic (1998) WHO Geneva
		Wilson T.M., de Lisle, G.W. and Collins, D.M. (1995) Effect of <i>inhA</i> and <i>katG</i> on isoniazid resistance and virulence of <i>Mycobacterium bovis</i> . <i>Mol. Microbiol.</i> 15, 1009 - 1015
		Yuan, Y., Crane, D.C., Simpson, R.M., Zhu, Y.Q., Hickey, M.J., Sherman, D.R. and Barry, C.E. III. (1998) The 16 kDa $\alpha$ -crystalline (Acr) protein of <i>Mycobacterium tuberculosis</i> is required for growth in macrophages. <i>Proc. Natl. Acad. Sci. USA</i> 95, 9578 - 9583
		Zahrt, T.C. and Deretic, V., (2000) An essential two-component signal transduction system in <i>Mycobacterium tuberculosis</i> . <i>J. Bacteriol.</i> 182, 3832 - 3838
		Zahrt, T.C and Deretic, V. (2001) <i>Mycobacterium tuberculosis</i> signal transduction system required for persistent infection. <i>Proc. Natl. Acad. Sci. (USA)</i> 98, 12706 - 11
		Zhang, Y. and Amzel, L.M. (2002) Tuberculosis drug targets. <i>Current Drug Targets</i> 3, 131 - 154

NY02:505269.1

6

Examiner



Date Considered

12-20-04

\* Examiner: Initial citation considered, whether or not citation is in conformance with MPEP 609; Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.